

Conference Volume

6th Symposium for Research in Protected Areas

2 – 3 NOVEMBER 2017 – FACULTY OF NATURAL SCIENCES, UNIVERSITY OF SALZBURG



SUPPORTED BY THE FEDERAL GOVERNMENT AND THE EUROPEAN UNION

BUNDESMINISTERIUM
FÜR NACHHALTIGKEIT
UND TOURISMUS



The European
Agricultural Fund for
Rural Development
Europe investing in
rural areas



 **Nationalpark[®]
Hohe Tauern**

NATIONAL
PARKS
AUSTRIA



Our natural heritage.



6th Symposium for Research in Protected Areas

2 to 3 November 2017, Faculty of Natural Sciences, University of Salzburg, Austria

Conference Volume

doi: https://doi.org/10.1553/np_symposium2017
ISBN-13 Online: 978-3-7001-8317-4

Weblink: https://epub.oeaw.ac.at/symposium_on_protected_areas

© by Salzburger Nationalparkfonds, Gerlos Straße 18/2, 5730 Mittersill, Austria, 2018

SUPPORTED BY THE FEDERAL GOVERNMENT AND THE EUROPEAN UNION

BUNDESMINISTERIUM
FÜR NACHHALTIGKEIT
UND TOURISMUS



The European
Agricultural Fund for
Rural Development:
Europe investing in
rural areas



Table of Contents

AIGNER, S. & EGGER, G.: An application example for modeling grazing intensity: National Park Hohe Tauern	1
ANDERWALD, P., MILLHÄUSLER, A., HAENI, M. & HALLER, R.M.: Rain, cost or publicity – what determines variation in visitor numbers to the Swiss National Park?	5
ANDRIČ, M., ŠMUC, A., OGRINC, N., RAPUC, W., SABATIERE, P., VON GRAFENSTEIN, U. & ARNAUD, F.: Palaeoecological Research in the Triglav National Park (Slovenia)	7
ARNBERGER, A., ALLEX, B., EDER, R., HUTTER, H.-P., WALLNER, P., BAUER, N., ZALLER, J.G. & FRANK, T.: Protected areas' landscapes as resources for human health and well-being – case studies from Austria	11
ARPIN, I., COSSON, A., MAZARD, C., RONSIN, G. & SCHEURER, T.: Collaboration between researchers and protected area managers – Empirical insights	15
AURENHAMMER, S., KOMPOSCH, C. & KREINER, D.: Where do endemics reside? Endemic beetles in the Gesäuse National Park (Austria) and their relevance for nature conservation	19
AZZONI, R.S., FUGAZZA, D., ZERBONI, A., SENESE, A., D'AGATA, C., MARAGNO, D., CARZANIGA, A., CERNUSCHI, M. & DIOLAUTI, G.A.: The evolution of debris mantling glaciers in the Stelvio Park (Italian Alps) over the time window 2003-2012 from high resolution remote-sensing data	25
BACKHAUS, N. & GRAEFE, O.: Park objectives in transformation: What are parks for and why do we want them?	29
BADURA, J. & FORSTER, S.: Parks as an instrument for sustainable regional development. A study in the intersection of tourism research, cultural studies and aesthetics	33
BANESCHI, I., GIAMBERINI, M.S., VITERBI, R., CERRATO, C., IMPERIO, S., FERRARIS, S. & PROVENZALE, A.: Changes in Alpine grassland of Grand Paradiso National Park (Italy): first results from CO ₂ fluxes monitoring programme	35
BARGMANN, T. & VETAAS, O.R.: The management of wild reindeer (<i>Rangifer tarandus</i>) in Hardangervidda National Park, Norway	39
BAUMGARTNER, M. & GOLLMANN, G.: Conservation of the yellow-bellied toad in Gesäuse National Park: collecting baseline data	41
BEIERKUHNLEIN, C. & IRL, S.D.H.: Threats of Climate Change to Single-Island Endemic Species in Protected Areas	45
BERNSTEINER, H., GÖTZ, J., SALCHER, B., OTTO, J.-C. & LANG, A.: Postglacial evolution and recent siltation of the protected lake Taferlklaussee (Salzkammergut, Upper Austria)	49
BETHWELL, C.: People in German Biosphere Reserves	55
BETHWELL, C., SATTLER, C., VAN BUSSEL, L., GIERSCH, G. & MEYER, A.: Can collaborative governance approaches enhance the spatio-temporal fit between agricultural related ecosystem services?	57
BILD, C. & OPP, C.: Nature Tourists and National Parks: How important are Nature and the Protection Status of an Area for Visitors?	59
BIONDA, R., RUBOLINI, D., ROTELLI, L. & IMPERIO, S.: Altitudinal shifts of Alpine grouse in the Veglia-Devero Natural Park, western Italian Alps	61
BLASCHKA, A., GUGGENBERGER, T. & HUBER, R.: Tough forage on alpine pastures? Nutritional value, phenology and stand structure of alpine pastures over 20 years	65
BLATTNER, L. & VON FUMETTI, S.: Faunistic assemblages of natural springs in different areas in the Swiss Nationalpark	69
BÖHM, J.: Basic principles for the optimization of wildlife management in 'Donau-Auen National Park'	73

BONDAR-KUNZE, E., KECKEIS, H., RÖSLER, S., GMEINER, P., LIEDERMANN, M. & HEIN, T.: How ship-induced wave trains affect shoreline communities in regulated rivers	75
BRANG, P.: Challenges for future wildlife management in and around protected areas	79
BRAUN, V. & HALLER, A.: Trajectories of Protected Area Creation in the Austrian Alps: two case studies	83
BRODT, M.S.C., BERGMANN, M., BROMAN, E.N., SANFELICE, G., FERREIRA, J.D., LUNARDI, L., HÜLLER, A. & DE CARLI, L.: Environmental damages in the Atlantic Forest Biome: A case study	85
BUCKEL, J., OTTO, J.-C., PRASICEK, G. & KEUSCHNIG, M.: The formation of glacial lakes in Austria and in the Hohe Tauern National Park since the Little Ice Age	89
BÜRGI, S. & WALLNER, A.: Increasing Research Interest in Protected Areas in Switzerland	93
CAZACU, C. & ADAMESCU, M.C.: Ecosystem services provided by the bio-physical structure of natural capital in the Danube Delta Biosphere Reserve Romania	97
CERRATO, C., VITERBI, R., BIONDA, R., VETTORAZZO, E., PEDROTTI, L., MOVALLI, C. & PROVENZALE, A.: A multi-taxa approach in mountain ecosystem: a shared protocol between 6 Italian Parks	103
CHURAKOVA (SIDOROVA), O.V., SAURER, M., BRYUKHANOVA, M., SIEGWOLF, R. & BIGLER, C.: Water-use strategies of conifer trees from the Swiss National Park to recent climatic changes	107
CLEMETSEN, M. & STOKKE, K.B.: Connecting nature, local cultures and tourism in subarctic landscape – a case study of local communities attached to the Varanger Peninsula National Park in Northern Norway	111
DEMETZ, A. & SCHULZE, C.H.: Effects of floodplain dynamics on richness, abundance, composition and functional diversity of grasshopper assemblages in the Donau-Auen National Park (Austria)	115
DIRNBÖCK, T., KOBLER, J., KRAUS, D., SCHINDLBACHER, A., SEIDL, R. & MIRTL, M.: From long-term ecosystem monitoring to regional modelling of ecosystem function in the National Park Kalkalpen, Austria	119
DUSCHER, T.: Invasion of non-indigenous carnivores in Austria – the current status of the raccoon (<i>Procyon lotor</i>) and the raccoon dog (<i>Nyctereutes procyonoides</i>) and prospects of their future distribution	123
DVORAK, M., BIERINGER, G., GRÜLL, A., KARNER-RANNER, E., KOHLER, B., LABER, J., NEMETH, E., RAUER, G. & WENDELIN, B.: Long term-monitoring of birds reveals drastic changes in the bird communities at the national park Neusiedler See - Seewinkel	125
EGGER, G., AIGNER, S. & POSCH, K.: Modeling grazing intensity of grassland	129
EIBES, P.M., KIENLE, D. & BEIERKUHNLEIN, C.: Endemic vascular plants in the high mountains of the Sierra Nevada National Park (Spain)	133
ELMAUER, K., CATTOEN, E.-M., LENZE, B. & BERGHAUSEN, K.: Influencing behaviour of visitors and residents in protected areas	137
FALLANCA, C., CARRÀ, N. & TACCONE, A.: Aspromonte National Park – The heart of the Metropolitan City of Reggio Calabria: design scenarios	141
FAVILLI, F., BÍL, M., SEDONÍK, J., ANDRÁŠIK, R., KASAL, P., AGREITER, A., SCHOBER, L., SICHER, P., GERSTGRASSER, L. & STREIFENEDER, T.: Combined use of KDE+ software and empirical observation to identify animal-vehicle collisions' hotspots in South Tirol, Northern Italy	145
FORTI, L., NOTARO, S. & PALETTO, A.: Sustainable hunting plan as a tool of wildlife management: the Italian case	151
FRITZ, L., RIEDE, M., KUTHE, A., KELLER, L. & STÖTTER, J.: <i>YOUrALPS</i> : Integrating young people's concepts, educators' teaching principles and stakeholders' educational tasks for a transformation of education and society towards sustainability in the Alps	157
FUNK, A. & HEIN, T.: Analysing the impact of drivers and pressures on the conservation goals of protected areas along a large navigable river, the Danube River	161
FÜREDER, L., NIEDRIST, G.H., SCHÜTZ, S.A.: Monitoring Alpine rivers: recent progress and future challenges	163
GABROVEC, M. & PAVŠEK, M.: The Triglav Glacier: Seventy Years of Regular Observations	169

GAHBAUER, C.: The National Park Gesäuse Partnership: From resistance to cooperation	173
GEDON, L., SONNENSCHNEIN, R. & WALZ, A.: Remote sensing based comprehensive monitoring of land cover change in protected areas	177
GIERSCH, G. & MEYER, A.: Tackling jointly the challenges of rural development and biodiversity conservation in protected areas	179
GRAF, R., AELVOET, P., GUYOT, C. & SPAAR, R.: Biology, conservation, and promotion of Skylark populations in subalpine and alpine zones of the Beverin-Naturpark, Switzerland	181
GRAFETSTÄTTER, C., GAISBERGER, M., PROSSEGGER, J., RITTER, M., KOLARŽ, P., PICHLER, C., THALHAMER, J. & HARTL, A.: Effects of green exercise and waterfall aerosol on mucosal immunity and chronic stress – A randomized controlled clinical trial	185
GREILINGER, M., SCHAUER, G. & KASPER-GIEBL, A.: Monitoring of Saharan Dust – Influence on aerosol composition and snow chemistry	189
GRISCHOTT, R., KOBER, F., LUPKER, M., REITNER, J.M., DRESCHER-SCHNEIDER, R., HAJDAS, I., CHRISTL, M. & WILLET, S.D.: Millennial scale variability of denudation rates for the last 15 kys inferred from the detrital ¹⁰ Be record of lake Stappitz in the Nationalpark Hohe Tauern, Austrian Alps	193
GRUBER, V.M.: How much wilderness is left in Europe's Wilderness Areas? A comparative analysis of selected wilderness areas in Central Europe	197
GUMPINGER, C.: The current situation of the River Enns fish fauna around the Gesäuse National Park	199
HADAR, L. & PEREVOLOTSKY, A.: Adaptive management at the Ramat Hanadiv Nature Park, Israel: Expectations vs. Reality in a dry Mediterranean ecosystem	201
HAELER, E., HINDENLANG, K., PELLISSIER, L. & LACHAT, T.: Biodiversity and forest structures at the Zurich Wilderness Park Sihlwald	205
HARTL, A.: Nature therapy and green exercise as remedies for emerging civilization diseases	207
HARTMEYER, I., KEUSCHNIG, M., FEGERL, L., VALENTIN, G., HELFRICHT, K. & OTTO, J.-C.: Long-term monitoring of climate-sensitive cirques in the Hohe Tauern range	209
HECKE, C., JUNGMEIER, M. & KREINER, D.: Patterns of Wilderness – en route to compiling an inventory of the national processes in Gesäuse National Park (Ennstaler Alpen)	213
HEDINGER, C. & FRIEDLI, A.: How can Swiss Regional Nature Parks improve the functionality of the ecological infrastructure? Pilot-scheme in two Nature Parks in the Canton of Berne	217
HEIN, T., BONDAR-KUNZE, E., FUNK, A., PÖLZ, E.-M., PITZL, B. & WEIGELHOFER, G.: Restoring fluvial landscapes – ecological effects of side-arm reconnection	221
HENNIG, S.: The use of crowdsourced (spatial) data in visitor management – Discussed by the example of a visitor hotspot in Berchtesgaden National Park	225
HERRMANN, E. & HÖBINGER, T.: Melting underground ice masses in Hochtor massif, Gesäuse National Park – documentation and implications for the water resources of alpine karst	229
HÖDL, C.: Geocaching in Austrian National Parks	235
HOFFMANN, S. & BEIERKUHNLEIN, C.: Evaluating the potential of protected areas to preserve biodiversity at large scales	237
HÖNIGSBERGER, B.: Wintering birds in floodplain forests – Effects of vegetation structure and landscape composition on species assemblages	239
HUMER-GRUBER, A.: Sustainable regional development approaches in Alpine biosphere reserves – Farmers' perceptions	243
HUMMEL, C., BOYER, Y., JUREK, M., ANDRESEN, P.M., KOBLE, J., BEIERKUHNLEIN, C., PROVENZALE, A., ZIV, G., HEURICH, M., KORDELAS, G., DE WIT, R., MANAKOS, I. & HUMMEL, H.: Ecosystem Services and Pressures in European Protected Areas: Divergent Views of Environmental Scientists and Managers	247
HUNZIKER, M. & HUBSCHMID, E.: Experimental Long-Term Evaluation of a Campaign to Reduce Freeriding-Wildlife Conflicts of Snow Sports	249
HYNEK, B., WEYSS, G., NEUREITER, A., BINDER, D. & SCHÖNER, W.: From annual glacier mass balances towards a remote monitoring of near real-time mass changes	251

IMPERIO, S., RANGHETTI, L., VON HARDENBERG, J., PROVENZALE, A., BONCOMPAGNI, E. & FASOLA, M.: Effects of protection status, climate, and water management of rice fields on long-term population dynamics of herons and egrets in north-western Italy	255
JAHIĆ, H., SPAHIĆ, M. & TEMIMOVIĆ, E.: Prokoško Lake – a monument to man	259
JUTZI, D.: Supply of local products in Parc Ela – constraints and possible solutions	265
KAHLEN, J., SVADLAK-GOMEZ, K., BEIGLBLÖCK, C. & WALZER, C.: How hunting and legal regulations shape ecological connectivity in the alpine region	267
KASPER-GIEBL, A.: The Lab Above the Clouds – Sonnblick Observatory and Nationalpark Hohe Tauern	269
KATZENSTEINER, K., DARABANT, A., MAYER, M., MATTHEWS, B., PRÖLL, G. & SCHAUFLENER, J.: Open Wounds in Thin Skin: Soil Processes after Natural Disturbances	273
KERLE, S. & TAPPEINER, U.: The Tyrolean Alps LTSER platform – connecting science and people	277
KEUSCHNIG, M. & HARTMEYER, I.: The Open Air Lab Kitzsteinhorn (OpAL) – Open Innovation in High Altitude	281
KHIL, M.-L. & SCHULZE, C.H.: Identifying crucial factors for nest survival and predation in a northern lapwing <i>Vanellus vanellus</i> population in the Lake Neusiedl - Seewinkel National Park	285
KIENLE, D., EIBES, P.M. & BEIERKUHNLEIN, C.: Survival in litte? Refugia of high-elevated plants in the Spanish Sierra Nevada	287
KLASZ, G., BAUMGARTNER, C. & GUTKNECHT, D.: An improved bedload management for the Danube River in the Donau-Auen National Park. An application of the ‘principle Sisyphus’	291
KLOSTERHUBER, R. & VACIK, H.: Naturalness and conservation status of forest habitats in the National Park Hohe Tauern Salzburg (Austria)	297
KNAPP, E. & SCHULZE, C.H.: Seasonal patterns of food use of wild boar (<i>Sus scrofa L.</i>) in a Central European floodplain forest	303
KNAUS, F.: Park-labelled products as a tool for innovation and regional development	307
KNOLL, T. & REITER, K.: Development of Hybrid Poplar Stands in the Donau-Auen National Park (Austria)	309
KÖCK, G.: Research in protected areas funded by the Austrian National Committee for UNESCO’s ‘Man and the Biosphere’ programme	313
KOHLER, B., PLUTZAR, C., ENZENHOFER, K., SCHRANK, J. & ZIKA, M.: Wilderness preserved? Representation of wild land within Austria’s network of protected areas	317
KOHLER, Y. & PLASSMANN, G.: Ecological connectivity in the Alps and beyond – a long term challenge	321
KOMPOSCH, C.: A new classification of endemic species of Austria for nature conservation issues	323
KÖPPL, W. & OBERKLAMMER, I.: <i>Dianthus plumarius</i> subsp. <i>blandus</i> – Distribution and Habitat Features	327
KÖRNER, C.: Comparative, long-term ecosystem monitoring across the Alps: Austrian Hohe Tauern National Park, South-Tyrol and the Swiss central Alps	331
KOZYRAKI, M., NYKTAS, P., KYRIAKOPOULOU, N. & PLOUMI, K.: Ecosystem Services: Contribution of Natura 2000 Network in the economy and society of Crete	339
KRUSE, A. & STRASSER, P.: A tourism toll road in the biggest central European National Park – a paradigm? Or an antagonism? And what about the inhabitants’ expectations? – An Example from the Hohe Tauern National Park	345
KUDRNOVSKY, H. & LAZOWSKI, W.: Fluvial landscapes – an important ecological dimension within European protected areas	349
KUPFERSCHMID, A.D.: Local trophic cascading impact of wolves on tree regeneration in summer and winter areas of ungulates	353
KÜRY, D., LUBINI, V. & STUCKI, P.: Index to evaluate the vulnerability to climate change of Mayfly, Stonefly and Caddisfly species in alpine springs	357
KUTTNER, M.: Biodiversity and Landscapes: Where is the missing link?	361

LAMPRECHT, A., PAULI, H., FERNÁNDEZ CALZADO, M.R., STEINBAUER, K. & WINKLER, M.: Unique and highly threatened – endemic plants at the cold edge of southern Europe	363
LEITNER-KLAUNZER, V.: Education of environmental consciousness as an education goal of the partnership between national park and new secondary school. Results of a qualitative evaluation study	367
LEONHARDT, G., SEIFERT, L., GERECKE, R., MÜLLER, J., HOTZY, R. & LOTZ, A.: Springs in the Bavarian National Parks as indicators for climate change	369
LEONHARDT, G., WARSCHER, M., STRASSER, U. & KUNSTMANN, H.: Water balance modeling and climate impact research in the Berchtesgaden National Park	373
LIEB, G.K., BAUER, C. & KELLERER-PIRKLBAUER, A.: Temporal and spatial variability of bedrock, debris and glaciers in the Austrian Alps since the Alpine Last Glacial Maximum and its relevance for ecological research	375
LIEB, G.K. & KELLERER-PIRKLBAUER, A.: Monitoring of abiotic natural processes in the Hohe Tauern National Park, Austria: A long-term approach	379
LIPP, S., OETTEL, J., STEINER, H. & FRANK, G.: Dynamic Processes in Austrian Natural Forest Reserves	383
LOTTA, F. & SCHILLECI, F.: The Sicilian system of Protected Areas – Integrated planning and management policies for protected areas	387
LOTZ, A. & ORAVEC, A.: Landscape Monitoring in Berchtesgaden National Park – Comparative spatio-temporal Analysis of Land Cover Inventories	391
LUTHARDT, V., PROBST, R. & LÜDICKE, T.: Design, implementation and performance of an ecosystem monitoring program in Biosphere Reserves – the long-term observation of ecosystems in the Biosphere Reserves in the federal state Brandenburg (ÖUB)	395
MANEA, G., TIȘCOVSCHI, A., VIJULIE, I., MATEI, E., CUCULICI, R., PEDA, M. & COCOȘ, O.: Interspecific relationships within Romania's protected areas. Case study: the cohabitation <i>Homo sapiens sapiens</i> - <i>Ursus arctos arctos</i> , in Harghita Mountains	401
MARGRAF, C.: Promoting terrestrial habitat- and species-diversity as consequence of river rewilding (middle Isar, Natura 2000-area)	405
MARINGER, A.: Biodiversity assessment in the Gesäuse National Park	409
MAYRHOFER, S., KIRCHMEIR, H., WEIGAND, E. & MAYRHOFER, E.: Assessment of forest wilderness in the Kalkalpen National Park	413
MBENG, E. & BUBA, U.H.: Community livelihoods versus biodiversity conservation. Constraints to sustainability of the Santchou forest reserve in the Western highland region of Cameroon. Protected areas' landscapes as resources for human health and wellbeing	417
MICEK, M.: Study on the introgression of hybrid poplar genes in the gene pool of black poplar in the Danube National Park	421
MICHEL, A.H.: 'What is a national park for?' – Principles of worth in a Swiss national park project	425
MILJIĆ, S., MAKSIN, M., KRUNIĆ, N. & RISTIĆ, V.: Zoning in spatial planning for protected areas and tourism destinations in mountain regions - case of the Kopaonik National Park in Serbia	429
MITROFANENKO, T., SNAJDR, J., MUHAR, A., PENKER, M. & SCHAUPPENLEHNER-KLOYBER, E.: Potential for applying Intergenerational Practice to involve underrepresented age groups in protected area management	433
MRAK, I., ODAR, M., MAROLT, M., KREK, A., BREZNIK, K. & HALVORSON, S.J.: Aspects of carrying capacities and recreation management: The case of Triglav National Park, Slovenia	437
NAGL, C. & SCHULZE, C.H.: Population density and habitat preferences in a Tawny Owl <i>Strix aluco</i> population in floodplain forests in Eastern Austria	441
NEUMÜLLER, U., PACHINGER, B. & FIEDLER, K.: Impact of inundation regime and meadow management on wild bee communities and bee-flower networks in the Donau-Auen National Park	445
NEWESELY, C. & TAPPEINER, U.: Climatological reference data of a newly established long-term monitoring program in the central Alps	449
NICKLAS, L., MALLAUN, M. & ERSCHBAMER, B.: Winners and losers of climate change in the Central Alps	453

NIEDRIST, G.H.: Spatial and temporal variation of chironomid assemblages in high altitude streams of the Hohe Tauern Nationalpark: environmental niche differentiation	455
NIEDRIST, G.H. & FÜREDER, L.: Feeding plasticity of alpine stream chironomids: evidence from river monitoring in the Hohe Tauern NP	457
NIGL, P.: The expansion of the Kalkalpen National Park from the perspective of political ecology	461
NOLL, D.: Small ruminants in a sustainable socio-ecological metabolism: A case study from Samothraki, Greece.	463
OBERHUBER, W.: Water availability as a key factor of forest dynamics in protected areas – long-term perspectives inferred from tree rings	465
OBERKLAMMER, I. & KÖPPL, W.: <i>Dianthus plumarius</i> subsp. <i>blandus</i> – Monitoring under extreme conditions	467
OETTEL, J., FRANK, G., STEINER, H. & LIPP, S.: Conflict Management – Case studies from the Austrian Natural Forest Reserve Programme	471
OLEFS, M., KOCH, R., HIEBL, J., HASLINGER, K. & SCHÖNER, W.: Seasonal snow cover evolution in the Nationalparks Austria since 1961	475
ORAVEC, A.: Habitat selection of alpine chamois under different climatic conditions in the Alpine and Carpathian mountain chains	479
OTTO, J.-C., HELFRICHT, K., PRASICEK, G., KEUSCHNIG, M., BINDER, D. & BUCKEL, J.: Predicting future glacial lakes in Austria – preliminary results	483
PALETO, A., DE MEO, I., DOBŠINSKÁ, Z., LAKTIĆ, T., MALOVRH, Š.P. & ŠÁLKA, J.: Stakeholders' opinions about the relevance of human activities in Natura 2000 sites: comparison among three European countries (Italy, Slovenia, Slovakia)	487
PAWLICZEK, P.: Bioclimatic indices in the context of biodiversity, Karkonosze/Giant Mts., SW Poland	493
PICHLER, C., SALLETMAIER, C., BAUERNBERGER, L., HAAS, W., LEHOFER, B., KUHN, W., BOTT, A., STECKENBAUER, C. & HARTL, A.: The Alpine Health Region Salzburg - an evidence-based health tourism approach for the valorization of natural resources within and outside protected areas	497
PICHLER-KOBAN, C., UKOWITZ, M. & JUNGMEIER, M.: Vertigo Effect: Institutional dynamics in nature conservation	501
PICHLER-SCHEDER, C. & GUMPINGER, C.: The recovery of ecologically and chemically impaired tributaries in the Podyjí/Thayatal National Park	505
PITTINO, F., FRANZETTI, A., NAVARRA, F., TAGLIAFERRI, I., GANDOLFI, I., BESTETTI, G., MINORA, U., AZZONI, R.S., DIOLAIUTI, G., SMIRAGLIA, C. & AMBROSINI, R.: Bacterial communities changes in cryoconite on an alpine glacier	509
PLANK, C., BOHNER, A. & WRBKA, T.: The effects of mountain farming on biodiversity-monitoring and evaluation of vegetation changes on managed and abandoned mountain pastures in the Gesäuse National Park (Styria, Austria) in an eleven years timescale	513
PÖLLINGER, U.: Are we willing and are we able to protect endemic species from becoming extinct?	517
PÖLZ, E.-M., BONDAR-KUNZE, E., WEIGELHOFER, G., ZHENG, X. & HEIN, T.: The importance of heterogeneous shoreline habitats for ecosystem functions in large regulated rivers	519
PROBONAS, M., PLOUMI, K., KYRIAKOPOULOU, N., XIROUCHAKIS, S., PILIGOTSI, G., GEORGOPOULOU, E. & BAXEVANI, K.: Ecosystem Services in NATURA 2000 areas of Crete: Information and communication campaign for the ecological, social and economic values of ecosystem services	523
REMPFLER, T.: Where do red deer come from and where do they go?	529
REMSCHAK, C.: Springs in Gesäuse National Park - Hotspots of biodiversity	531
RESCH, S. & BLATT, C.: Assessing small mammal community diversity with minimally invasive field methods – examples from the Nationalpark Gesäuse (Austria)	537
RESL, F.: Geomorphological Trails in Austria	541
RIEDL, C. & REISENHOFER, S.: Monitoring Permafrost at Hoher Sonnblick, Hohe Tauern, Austria	543

RIEDLER, B. & LANG, S.: Using an aggregated remote sensing-based habitat quality index for the identification of spatially targeted conservation measures	545
RIXEN, C.: Long-term changes in alpine tundra vegetation: 25 years of the International Tundra Experiment ITEX	549
ROBSON, B.A. & HÖBLING, D.: Automatic Glacier Monitoring in the Hohe Tauern National Park, Austria	551
ROSSI, C., RISCH, A.C., SCHÜTZ, M., KNEUBÜHLER, M., SCHAEPMAN, M.E. & HALLER, R.M.: A regional assessment of functional diversity in heterogenous grassland with different agricultural management	555
RUMPF, S.B., HÜLBER, K., ZIMMERMANN, N.E. & DULLINGER, S.: Climate-driven range dynamics and potential current disequilibrium in Alpine vegetation	559
RUOSS, E. & ALFARÈ, L.T.: Shifting protected area strategies to evidence based governance and management	561
SACHSER, F., NOPP-MAYR, U. & GRATZER, G.: Linking habitat dynamics and population cycles of small mammals in different mountainous forest types in Austria	565
SATTLER, C.: Let's cooperate: How collaborative governance approaches can help to address institutional fit in protection areas	567
SAUKEL, L.: Visitors' use of national park services and their affinity to the Gesäuse National Park	571
SCHABETSBERGER, R., JERSABEK, C., LUGER, M. & KREINER, D.: Contrasting top down effects of amphibians and stocked fish in Austrian alpine lakes	575
SCHAUFLER, J.: Soil succession in relation to vegetation on a subalpine forest fire site in the Northern Limestone Alps	577
SCHEURER, T.: Managing externally caused impacts on protected ecosystems in a long-term perspective – lessons learnt from the Swiss National Park	579
SCHIRPKE, U. & SCOLOZZI, R.: Participatory modelling for understanding consequences of management choices on ecosystem services and biodiversity in protected areas	581
SCHÖPFER, A.: Capturing the effects of the Danube River incision on the potential natural vegetation of the Donau-Auen National Park	585
SCHÖTTL, S.: The potential of UAV-data for surveying sediment dynamics - A case study in the Gesäuse National Park	589
SCHÜTZ, S.A. & FÜREDER, L.: Diversity measures indicating environmental change in alpine river ecosystems	591
SCHÜTZ, S.A. & FÜREDER, L.: Unexpected larval development in a highly glaciated headwater	595
SCHWIENBACHER, M.: Anthropogenic influence on primary succession: A comparative study of 3 glacier forelands of the Central -Alps, Austria	599
SEISER, B., FISCHER, A. & WIESENEGGER, H.: Five Years of Glaciological Monitoring of Venedigerkees, Hohe Tauern National Park, Austria	603
SEMERNYA, L., ANDRESEN, P.M., ALFTHAN, B., JUREK, M. & ADLER, C.: Waste management in mountain protected areas	607
SIEGRIST, D.: Whatsalp Wien – Nizza 2017. Eine Fußreise durch die Alpen und ihre Großschutzgebiete	611
SLOTTA-BACHMAYR, L.: Large predators in protected areas – risk or chance	613
STACHOW, U. & BETHWELL, C.: Land use in German Biosphere Reserves and its ecological impact	615
STADEL, C.: Interactions, challenges, and management issues at the fringe of National Parks: The case of the Riding Mountain Biosphere Reserve	617
STEINBAUER, K., LAMPRECHT, A., WINKLER, M., BARDY-DURCHHALTER, M., KREINER, D., SUEN, M. & PAULA, H.: Shifting composition and functioning in alpine plant communities – Evidence of climate warming effects from 14 years biodiversity observation in the Northeastern Alps	621
STEINER, H.: Detecting a change in diversity of vascular plants in the natural forest reserve 'Gaisberg' – a comparison of a systematic and representative sample approach	623

STOCKER-WALDHUBER, M. & FISCHER, A.: 10 years of glacier mass balance monitoring on Mullwitzkees (Hohe Tauern)	627
STÖHR, O.: Ferns in the spray: the pteridophyte flora of the Krimml Waterfalls	629
STOCKE, K.B. & CLEMETSEN, M.: Integrated planning of national parks and adjacent areas – possibilities and limits in cooperation for nature-based tourism and	633
STUMVOLL, M.J., GÖTZ, J. & BUCKEL, J.W.: Surface change modelling of small scale debris flow dynamics (Mühlsturzsgraben, National Park Berchtesgaden, Germany)	637
ŠVAJDA, J., MEESSEN, H. & KOHLER, T.: Participatory Management of Protected Areas in Slovakia: Reconciling nature conservation and local development	643
TEMIMOVIĆ, E., SPAHIĆ, M. & JAHIĆ, H.: Mountain lakes in National Park 'Sutjeska' – Evolutionary self-development	649
TESTER, U.: Indicators for good management of protected areas	653
THOM, D.: Disturbance impacts on forest succession, biodiversity, and ecosystem services in a changing world	657
VACIK, H., STEINER, H., FRANK, G. & RUPRECHT, H.: Long term monitoring of natural regeneration in natural forest reserves in Austria	659
VACIK, H., STEININGER, B., STEINER, H. & FRANK, G.: A Fagetum in the natural forest reserve Luxensteinwand - a rare element in the Waldviertel region	665
VANCURA, V.: Wilderness is a unique laboratory – Science and Wilderness in Europe	669
VIJULIE, I., TİRLÄ, L., MANEA, G., MATEI, E. & CUCULICI, R.: Old-growth forests of Domogled-Valea Cernei National Park (Romania) between conservation and mismanagement	675
VITERBI, R., CERRATO, C., BIONDA, R., BASSANO, B. & PROVENZALE, A.: Biodiversity monitoring in the NW Italian Alps: state and expected changes	679
VOLKMER, J.: The land snail fauna of the National Park Gesäuse – Ecology of the alpine land snails, with a special focus on endemic species	683
VON FUMETTI, S.: Ecology of springs in the Swiss National Park: first results and future plans	685
WAGNER, T., WINKLER, G., RIBIS, M., KELLERER-PIRKLBAUER, A., LIEB, G.K. & KRAINER, K.: Rock glaciers – prominent landforms in (protected areas of) Austria	689
WAHBI, A., AVERY, W.A., FRANZ, T.E., DERCON, G., HENG, L. & STRAUSS, P.: Mobile Soil Moisture Sensing in High Elevations: Applications of the Cosmic Ray Neutron Sensor Technique in Heterogeneous Terrain	693
WALZ, A. & KORUP, O.: CORINE for large-scale monitoring of PAs in Europe	697
WANZENBÖCK, J.: Endemism below the species rank: Population genetics of the European Mudminnow (<i>Umbra krameri</i>)	701
WARCHALSKA-TROLL, A.: National parks and Natura 2000 sites in Polish Carpathians vs local people: changing attitudes within the past 10 years	703
WARINGER, B.M., REITER, K., SCHULZE, C.H.: Population density and habitat preferences of the Collared Flycatcher (<i>Ficedula albicollis</i> Temminck, 1815) in floodplain forests – A case study from the Donau-Auen National Park, Lower Austria	707
WEINKE, E., HÖLBLING, D., ALBRECHT, F. & FRIEDL, B.: Interactive web services for landslide and habitat monitoring	711
WENDL, S.: The Social Construction of Nature, an explorative investigation of the constructed meaning of Nature within four national park exhibitions in Austria and Germany	715
WIDERIN, K. & REITER, G.: Bat activity above 3000 m in the Austrian Alps (Hoher Sonnblick, 3106 m)	719
WIESENEGGER, H., KUM, G. & SLUPETZKY, H.: 'Unterer Eisbodensee' – a good example for the future evolution of glacial lakes in Austria?	721
WILD, N.: Spatio-temporal patterns of dragonfly occurrence on meadows in the Donau-Auen National Park, Lower Austria	727

WILKES-ALLEMANN, J., LIEBERHERR, E. & BAERLOCHER, B.: Governing peri-urban forestry: filling the regulation gap with Swiss ‘Nature-discovery-parks’?	733
WINKLER, G., WAGNER, T., RIBIS, M., PAURITSCH, M. & KRAINER, K.: The impact of rock glaciers on the runoff of alpine catchments in protected areas of Austria	735
WINTER, M.-B., BAIER, R., MÜLLER, J. & AMMER, C.: Natural forest dynamics following bark beetle outbreaks in the Berchtesgaden National Park – Forest structure and biodiversity during disturbance and succession	739
WIPF, S. & RIXEN, C.: Long-term changes in summit plant diversity in the Swiss National Park	741
WITHALM, G., GETZNER, M. & KOHLER, Y.: Ecological Connectivity and its Contribution to a Green Economy	745
WITTMER, G.: Record and description of the sediments of the flood in June 2013 in the area of the Donau-Auen National Park	749
WOLF, A. & VOGEL, B.: Speleology in the Berchtesgaden National Park Eiskapelle: 26 years of surveying	753
WUNDER, A.: Comparative population genetic analysis of brown trout (<i>Salmo trutta</i>) from Kalkalpen National Park	757
WURSTER, M.T., WEIGELHOFER, G., PICHLER-SCHEDER, C., HEIN, T. & PÖPPL, R.: Sedimentological downstream effects of dam failure and the role of sediment connectivity: a case study from the Bohemian Massif, Austria	759
ZAGEL, B., SLUPETZKY, H. & WIESENEGGER, H.: The mass balance series of Stubacher Sonnblickkees 1946–2016 and the semi-direct calculation of the mass balance of glaciers. A contribution to LTER Austria	761
ZHENG, L. & PEER, T.: Disturbance and recovery of Biological Soil Crusts (BSCs) in the high alpine region of the Hochtorn (Grossglockner, Austria)	765
ZIMMERMANN, T. & KREINER, D.: ArcGIS-generated map of FFH-habitat types for Natura-2000 site Ennstaler Alpen/Gesäuse (Styria, Austria)	769
ZITTRA, C., VITECEK, S., UEBLEIS, S., WARINGER, J. & FUEHRER, H.-P.: Benefits of mosquito surveillance programs in protected areas: two case reports from Eastern Austria	773
ZOLLNER, D., FALKNER, J., EGNER, H. & JUNGMEIER, M.: Best practice Science_Linknockberge – Benefits and challenges of five years of co-operation between biosphere reserve and university	777
ZULKA, K.P., BIERINGER, G. & WEIGAND, E.: What is biodiversity, how can it be quantified and prioritised and what does that mean for Austrian national parks?	781

Tough forage on alpine pastures? Nutritional value, phenology and stand structure of alpine pastures over 20 years

Albin Blaschka, Thomas Guggenberger, Reinhard Huber

Abstract

Within the project presented, running from 2015-2018, nutritional value, phenology and the development dynamics of 16 alpine pastures in or close to the national park Gesäuse (Styria, Austria) are studied. Based on data from the years 1993-1996, ripening stages and development of raw fibre content of forage harvested are compared. Together with weather data, conclusions on temporal changes of energy content are drawn and guidance for good pasturing (stocking rate, start time of pasturing, duration) in the face of climatic changes can be given, also in regard to animal welfare. For the analysis, not the macroscopic phenological stages are used, but the raw fibre content and nutritional value/energy content, which can be analyzed more exactly and describes as a sum parameter quite well local growing conditions for plants.

As the project is still running at the time of the writing, all data and conclusions are only preliminary, but give already a coherent picture.

Keywords

alpine pastures, phenology, fodder value, adaption to climatic changes, land management

Introduction

Climate parameters are a proxy for several site parameters and represent abiotic site conditions, which influence directly the local vegetation. Ongoing and expected changes in climatic conditions will cause changes both on a species and community level, especially at higher elevations (KROMP-KOLB et al. 2014). Such changes in the metabolism of plants can be monitored by their metabolites, which influence the nutritional value for animals, among other properties (ABDELGAWAD et al. 2014). In our context, the most important relationship is that the phenologic age of a plant correlates positively with the raw fibre content.

These changes have an immediate effect on the suitability of high alpine pastures with its primary use of vegetation as forage for ruminants. A well-adapted use of the resources is the basis for a sound, sustainable agriculture which is the key for our cultural landscape, protected or not.

Therefore, based on the literature, the following hypotheses were formulated as the basis for this study:

- Hot summer cause an early start of vegetation period and accelerates the development of the plants and leads to an early ripening
- This accelerated development leads to a higher content of raw fibre in plants at the end of the vegetation period, which gives a decreased nutritional value.

Based on the hypotheses, we try to answer the following research questions:

- How high is the raw fibre content and the resulting nutritional value over the years (1993-1996 and 2015-2018) on selected plots?
- How does the within-season development of raw fibre content look like?
- Are there changes in the altitudinal gradient of raw fibre content between the two study periods (1993-1996 versus 2015-2018)?

Methods

Between 1993 and 1996 factors influencing yield and forage quality were studied by our research institution (GRUBER et al. 1998). To be able to characterise changes, a similar project was started in 2016 on the exact same locations, preceded by a small pilot study in 2015.

The study area is a 16km long transect north-east orientated, between the 'Schoberpass' and the town of Hieflau in the Gesäuse (see map in Fig. 1). The study area is in Styria, Austria, halfway between Liezen and Leoben (47.494°N and 14.668°E).

The line of the transect, which was defined already for the project 1993 – 1996 ('baseline'), cuts a mountain range with siliceous bedrock in the southern half and a range with calcareous bedrock in the northern half. Plots were fenced exactly at the following altitudes: 1100, 1300, 1500 and 1700. Crossing two mountain ranges, this gives 16 plots, 8 on each range, 4 oriented south, 4 oriented north (see Fig. 2).




Figure 1: Location of the study area around the national park Gesäuse, Styria, Austria.




Figure 2: The transect with the distributions of plots over the two mountain ranges

During the baseline project, the plots were larger than the re-installed ones in 2016, which have now an extent of 7.5 x 7.5 m and are fenced with barbed wire. A sample is taken from an area of around 7.5 x 2.5 m (ca. 18m²). During the pilot study 2015, no fixed fences were built; just one sample on each plot was taken.

For the baseline, the main sampling date for each altitude was defined as the beginning of flowering of the characteristic grass species. To be able to monitor the development two weeks before and two weeks after this main date, additional samples were taken, overall three samples per plot. The dates were documented as the respective day of the year. The sampling starts around mid-May on the plots at 1100m and ends mid-August on the plots at 1700 m altitude.

In the current project, the forage samples are taken on the exact same days of the year (+/- 2d) on the exact same locations as during the baseline project. A botanical assessment is done just before the second sampling within the plot (on around 35m²). This gives in total 48 forage samples and 16 botanical assessments per year.

The forage samples are analysed following the method of Weender for structural substances (see KIRCHGESSNER 2011) and minerals at our research institution. The samples for the year 2017 are in the lab for analysis at the time of the writing.

Results

The raw fibre content shows relatively high amplitude and increases only marginally with altitude. Due to the small sample size and the high variation, the results split by altitude and year are not statistically significant (see Fig. 4). But together with temperature data (not shown) a clear trend between years becomes visible. Hot years like 1994, 2015 and 2016, the raw fibre content is higher than in the other years. The difference between 1994 and 2015 is at +19 g raw fibre/kg dry matter, which equals a faster development of one phenological stage (shooting stage versus start of flowering) at the same time of the year (Fig. 3) which equals to a clear loss of nutritional value as the digestibility increases.




Figure 3: Raw fibre content by year. Hot years like 1994 and 2015 show a significantly higher content than the other years.




Figure 4: Raw fibre content by year and altitude. The differences are more complex, and due to small sample sizes no statistical significant effect can be shown.

Discussion

As the project is still running at the time of the writing, all data and conclusions are only preliminary, but give already a coherent picture.

The results clearly show a negative development of nutritional value of forage on alpine pastures in the study area during hot summers. What was found in 2015 and partly in 2016 was there already 1994, which was also one of the hottest years until then. The question if there are more hot summers than in earlier times was not part of this study. A hot summer leads to an earlier ripening of the forage available, by up to around two weeks. The longer vegetation period brings no advantages, as no fresh growth is coming. This is a challenge for all ruminants on alpine pastures, as with less energy the sometimes rough conditions on high pastures pose a serious threat, which touches also animal welfare questions. If there is a hot year, the altitudinal gradient in raw fibre content is also mostly flat, meaning there is a higher increase and bigger effect on the lower lying plots (1100 and 1300 msm) than on the higher ones (see Fig. 4).

Conclusion

In the course of history and the resulting traditions connected to alpine pasturing, dates for turn out and putting animals on pastures at different altitudes developed and proved useful, being site adapted and helping to avoid over use and damage to the pastures. These traditions get under pressure, be it because of a changing climate (like shown with our results) or other recent developments in the agricultural practice.

It is time for an evaluation of those traditional processes and management systems. Only with well adapted practices a sustainable, site adapted extensive land use will be possible in the future, which also helps to protect the co-developed biodiversity of our cultural landscape.

References

- ABDELGAWAD, H., PESHEV, D., ZINTA, G., VAN DEN ENDE, W., JANSSENS, I. A., & ASARD, H. 2014. Climate Extreme Effects on the Chemical Composition of Temperate Grassland Species under Ambient and Elevated CO₂: A Comparison of Fructan and Non-Fructan Accumulators. *PLoS One* 9 (3): 1–13. doi:10.1371/journal.pone.0092044.
- GRUBER, L., GUGGENBERGER, T., STEINWIDDER, A., SCHAUER, A., HÄUSLER, J., STEINWENDER, R., & SOBOTIK, M. 1998. Ertrag und Futterqualität von Almfutter des Höhenprofils Johnsbach in Abhängigkeit von den Standortfaktoren. In: Tagungsband 4. Alpenländisches Expertenforum „Zeitgemäße Almbewirtschaftung sowie Bewertung von Almflächen und Waldweiden“, 24. Und 25. März 1998. Irdning, Österreich: BAL Gumpenstein.
- KIRCHGESSNER, M., ROTH, F. X., SCHWARZ, F. J. & STANGL, G. I. 2011. Tierernährung. 13. Auflag. DLG-Verlag.
- KROMP-KOLB, H., NAKICENOVIC, N., STEININGER, K., GOBIET, A., FORMAYER, H., KÖPPL, A., PRETTENTHALER, F., STÖTTER, J. & SCHNEIDER, J. (Eds.). 2014. Austrian Panel on Climate Change - Sachstandsbericht Klimawandel. Verlag der Österreichischen Akademie der Wissenschaften, Wien, Österreich

Contact

Albin Blaschka
albin.blaschka@rauberg-gumpenstein.at
Höhere Bundeslehr- und Forschungsanstalt für Landwirtschaft Raumberg-Gumpenstein
Raumberg 38
8952 Irdning-Donnersbachtal
Austria
Phone: +43368222451-244